

## Review Article

Advancements in Life Sciences — International Quarterly Journal of Biological Sciences

ARTICLE INFO

Date Received: 30/07/2023: Date Revised: 31/08/2023; Date Published Online: 31/12/2023;

#### Authors' Affiliation:

Authors' Affiliation:

1. Department of Pathology, School of Medicine, Lorestan University of Medicine Sciences, Khorramabad - Iran

2. Razi Herbal Medicines Research Center, Lorestan University of Medical Sciences, Khorramabad - Iran

5. Symbiosis Institute of Business Management, Symbiosis Institute of Business Management, Symbiosis Institute of Business Management, Symbiosis Institute of Business (A. Baghada College of Medical Sciences, Department of Pharmacy, Baghada - Iraq

5. Gollege of Pharmacy, Al-Bayan

University, Baghada - Iraq

6. Department of Medical Laboratory Sciences, College of Applied Medical

7. Medical Laboratory Technology
Department, College of Medical

7. Medical Laboratory Technology
Department, College of Medical

Technology, The Islamic University, Najaf-

8. Scientific Research Center, Al-Aye Medicine, Shahid Beheshti University o Medical Sciences, Tehran - Irar Food Safety Division, Faculty of Public Health, Tehran University of Medical Sciences, Tehran - Irar

#### \*Corresponding Author:

Nasrollah Naghdi Email:

dr.naghdi93@gmail.com

#### How to Cite:

Adeli OA, Prasad KDV, Almajidi YQ, Khalaf HA, Hiazi A (2023). Phytotherapy in sexual disorder: overview of the most important medicinal plants effective on sexual disorders, Adv. Life Sci. 10(4): 505-514.

#### Keywords:

Diet quality; Dietary fiber; Food law: Food safety: Functional foods; Healthy eating

## Open Access





# Phytotherapy in sexual disorder: overview of the most important medicinal plants effective on sexual disorders

Omid-Ali Adeli<sup>1,2</sup>, KDV Prasad<sup>3</sup>, Yasir Qasim Almajidi<sup>4</sup>, Hasan Alauldeen Khalaf<sup>5</sup>, Ahmed Hjazi<sup>6</sup>, Beneen M. Hussien<sup>7</sup>, Hussein Ali Hussein<sup>8</sup>, Nasrollah Naghdi<sup>9\*</sup>, Mohadeseh Pirhadi<sup>10</sup>

## Abstract

lacksquare exual health is the result of the interaction of vascular, nervous and hormonal factors and is influenced by individual factors, interpersonal relationships, traditions governing family and society, culture and religion; these sexual disorders are one of the factors influencing marital life and it faces a challenge. The present study is a review of the most important medicinal plants effective on sexual disorders. In this study, a review of the key words "sexual function, sexual dysfunction, plant medicine, herbal medicine, treatment, fertility" was searched from Embase, Magiran, SID, Web of Science, Scopus, PubMed and Google Scholar databases. For each herb, selected published clinical trial and review articles were used. Based on the obtained results, medicinal plants such as Tribulus terrestris, Pomegranate, Red clover Lavandula angustifolia, Elaeagnus angustifolia, Pelargonium, Melissa officinalis, Quercus brantii, Ginkgo biloba, Trigonella foenum-graecum, Crocus Sativus, Zingiber officinalis, Ros Damascena, Celery seed, Date, Fennel and Carrot seed in clinical trials and Animal models are used to treat sexual disorders, Medicinal plants can effectively treat sexual disorders. Since hormone therapy has many side effects, increasing awareness of how herbs work on androgen, estrogen, and progesterone hormones and the sexual function of people will give us the knowledge that in couples' sexual dysfunctions, there are more choices for correction without using to have hormones with the aim of increasing the quality of life.

## Introduction

Marriage can be considered one of the most important and stable interpersonal relationships. However, depending on the quality of this relationship, marriage can have problematic consequences [1]. Marital satisfaction is one of the common concepts used to evaluate happiness and stability in marriage. In fact, what is more important than the marriage itself is the success of the marriage and the satisfaction of the couple [2]. Hawkins defined marital satisfaction as the feeling of satisfaction and happiness that a husband or wife experiences considering all aspects of their marriage; it is also one of the most important indicators of life satisfaction and family functioning [2]. One of the destructive factors of this satisfaction is sexual disorders, which cause emotional and psychological damage, decrease the quality and satisfaction of marriage, and increase marital exhaustion.

Therefore, any disorder that causes dissatisfaction with sex can lead to sexual dysfunction. dysfunction is defined as a person's inability to perform one or all phases of sexual activity, including desire, arousal, orgasm, erection/ejaculation for men, and lack of vaginal moisture for women [3]. They can cause sexual dysfunction as long as they affect any part of the sexual response cycle. There is limited information about the incidence and prevalence of female sexual dysfunction (FSD) in women. The incidence of FSD is estimated from 25.8% to 91.0% [4]. Decreased libido has the highest prevalence among lifetime sexual disorders for both men and women [3]. In a 5-year study of Finnish women aged 18-74 years, decreased libido was found in 45% of women: 20% among those younger than 25 years and 70-80% among those aged 55-74 years [4]. A 2006 survey by Hayes and his colleagues reported the prevalence of 16 problems in 64% of women, arousal problems in 31%, orgasm problems in 35%, and sexual pain in 26% [5]. Various researches also indicate the high prevalence of sexual disorders in Iranian women. The prevalence of sexual dysfunction in the general population of Iran is estimated at 35%, which is the highest prevalence among all sexual dysfunctions [3].

Endocrine disorders including breast and ovarian cancer, inflammatory diseases such as fibromyalgia and rheumatoid arthritis and neurological conditions such as multiple sclerosis, secondary (acquired) problems such as childbirth, hormonal changes, menopause, breastfeeding, trauma, psychological factors such as depression and anxiety, medications such as antidepressants, antipsychotics, and hormonal methods of birth control, lifestyle factors, such as unhealthy diet and weight, lack of exercise, smoking, and alcohol consumption, and psychosocial factors

such as age, education, income, and ethnicity can play a role in this [4, 6].

A recent large survey study in the United Kingdom, using proxy criteria for DSM-5 sexual disorders, found that while 22.8% of women reported one or more sexual problems including problematic orgasm, low sexual interest, and arousal or painful intercourse. There are now three criteria for diagnosing sexual disorders: symptoms must persist for at least 6 months, be experienced in all or nearly all (75% to 100%) sexual encounters or be persistent/recurrent, and clinically established [7]. In addition to women, men may also have sexual dysfunction. According to statistics, 52% of men aged 40 to 70 suffer from various degrees of sexual dysfunction caused by various biological and psychological factors. In addition, epidemiological evidence suggests that impotence affects 10 million American men [8]. Due to the sensitivity of the male reproductive system, it is easily affected by several risk factors such as chronic diseases, environmental pollutants, drug toxicity and unhealthy lifestyle, etc.

Maintaining normal sexual function depends on the coordination of several human systems, including the coordination of the nervous system, cardiovascular system, endocrine system, and reproductive system (9). A variety of factors such as psychological factors, surgery and trauma, vascular diseases, neurological factors, lifestyle changes, stressful living conditions, various chronic diseases, various environmental pollutants and drug side effects, etc.[9, 10]. The cause of FSD is multifactorial and has biological and psychological elements. In addition, many patients report concerns about different symptom clusters. Many women experience lack of sex or sexual problems in having an orgasm. When a woman is uncomfortable or unhappy with her feminine health, a woman has sexual dysfunction, which is called FSD. Diabetes, heart disease, cancer, arthritis, multiple sclerosis or alcohol abuse, high blood pressure medications, depression, oral contraceptives, endometriosis, cystitis, pelvic muscle problems, or chronic pelvic pain are causes the FSD [10]. Due to the multifactorial etiology of FSD, both pharmacological and non-pharmacological strategies have been used [11]. Although the sexual functions of men and women are affected by many aspects, currently there are almost no effective treatments in sexual medicine specifically for such disorders.

In the 1990s, sildenafil (brand name Viagra) was the first approved treatment for impotence to gain public attention. Chemical drugs, in addition to psychotherapy and cognitive therapy, are among the options for affected people. Despite the increase in access to effective modern medical treatments, the prevalence of side effects associated with these drugs

has led to the introduction of complementary medicine-based treatments as a desirable acceptable option in the treatment of various diseases as well as improving sex life [12]. Currently, although there are many traditional medicines or herbal medicines that have been used by patients with sexual dysfunctions with varying degrees of success, herbal products are still used irregularly in many regions and countries, and a large proportion of medicinal plants are They are traditionally used to solve sexual problems [9]. Iran is one of the first countries in using plants to treat diseases. The therapeutic effects of some plants have been investigated and proven by conducting scientific studies [13]. One of the methods of complementary medicine that can replace diets and medication is the use of medicinal plants. The use of medicinal plants, in addition to being economical, easy to access and low cost, have fewer side effects than chemical drugs.

The uses of medicinal plants provides a good reason to conduct new research in order to obtain information about the effect of medicinal plants in the treatment of problems and diseases [14]. Therefore, the present study is a review of the most important medicinal plants effective on sexual disorders.

## Methods

### Literature Search and Selection Criteria

In this review study, Embase, Magiran, SID, Web of Science, Scopus, PubMed and Google Scholar databases were searched using the keywords "sexual function, sexual dysfunction, plant medicine, herbal medicine, treatment, fertility". For each herb, selected published clinical trial and review articles were used. Low relevance or irrelevant articles, articles with irrelevant information, abstracts were the exclusion criteria.

## Discussion

In the present study, 21 species of plants were found that are used in the treatment of sexual disorders. The plants that are used to treat these disorders and their effectiveness are given in table 1.

#### Ros Damascena

It is especially useful for people who suffer from depression and have sexual disorders [30]. This plant has been especially effective in reducing dyspareunia and increasing orgasm and libido in depressed women [30]. In addition, the use of rose oil extract increases testosterone levels in men [31]. These functions are through the effect on sex hormones [30, 31].

#### Zingiber officinalis

It has improved the sexual performance of depressed people, especially in the area of arousal and desire in these patients [30] . In traditional medicine, it is used to improve sexual performance [30] in the study of Stein et al. (2018) which was conducted on 14 healthy men, 100 mg of the alcoholic extract of this plant improved erectile dysfunction and increased satisfaction in men, even middle-aged and elderly people [29] . Ginger has anti-inflammatory and antioxidant properties and reduces oxidative stress, thus helping to treat erectile dysfunction.

#### Crocus sativus

As an important factor related to the effect on nitric oxide and opioid secretion system, saffron has an efficient role on erection [27, 28, 34, 35].

#### Trigonella foenum-graecum

It is able to improve the overall score of sexual function in women [26, 36]. It has beneficial effects especially on the areas of arousal and sexual desire as well as dyspareunia [26, 37]. The results of a study showed that on 60 postmenopausal women, a group of 30 people received Trigonella foenum-graecum in the form of 5% vaginal cream for 8 weeks, there was a decrease in dyspareunia and an increase in libido in the Trigonella foenum-graecum group compared to placebo [26]. Trigonella foenum-graecum increases testosterone, which is a justification for increasing sexual desire (in both sexes). The increase in testosterone is normal and is related to the decrease in SHBG levels [36-40].

#### Cinnamomum verum (Cinnamon)

It is used to improve sexual performance, especially in the area of sexual desire and arousal [30, 41]. *Cinnamon* increases the humidity of the genital tract and as a result reduces dyspareunia and increases orgasm [30, 42].

#### Tribulus terrestris

It has been used for sexual disorders since ancient times. Studies have shown that consumption of *Tribulus terrestris* can increase sperm production and testosterone levels [43, 44]. This plant improves sexual performance in women especially by promoting sexual desire (especially during menopause) [13, 36].

The study by D'Souza et al. (2016) on 36 postmenopausal women who were treated with this plant for 120 days with 750 mg per day and in the placebo group, in all sexual areas except for moisturizing the vaginal environment, this plant improved the sexual performance of women. It also increased testosterone levels [13]. has been used for sexual disorders since ancient times. Studies have shown that consumption of *Tribulus terrestris* can increase sperm production and testosterone levels [43-46].

| Plant  | Name   | Type of study     | The number of participants | Target group                | Drug dosage and<br>drug form  | Duration of<br>intervention        | Result  | Main compound                                   |
|--|--|-------------------|----------------------------|-----------------------------|---|------------------------------------|---|---|
| Red Clover /<br>Khayatan et al.<br>(2019) [15]                     | Red clover                                       | Clinical<br>trial | 78                         | menopausal<br>women         | A 5 gram applicator   | 8 weeks                            | Improvement in all areas  | Trifolin  |
| Pelargonium /<br>Malkoti et al. (2016)                             | Pelargonium                                      | Clinical<br>trial | 120                        | menopausal<br>women         | Inhalation 3 times a<br>day, placing 2-3<br>drops of solution on<br>the skin of the<br>forearm  | 6 weeks                            | Improving sexual<br>performance in all<br>areas except<br>dyspareunia   | Pelargonium sidoides                            |
| Tribulus terrestris /<br>D'Souza et al. (2016)<br>[13]             | Tribulus<br>terrestris                           | Clinical<br>trial | 36                         | menopausal<br>women         | 750 mg (250 mg<br>tablet 3 times a day)   | 120 days                           | Improving all areas<br>except for moisturizing<br>the vaginal<br>environment and<br>increasing testosterone   | beta-D-<br>Fructofuranosyl                      |
| Granatum Punica L<br>/ Mohammadzadeh<br>et al. (2019) [14]         | Granatum<br>Punica L                             | Clinical<br>trial | 110                        | Non-<br>menopausal<br>women | Vaginal gel based<br>on Carbopol with a<br>concentration of 3%  | 8 weeks                            | Effective orgasm with increased frequency and increased sexual satisfaction   | Citric acid                                     |
| Quercus / Larzadeh<br>et al. (2016) [24]                           | Quercus  | Clinical<br>trial | 120                        | women                       | after the end of the<br>period for 3 days<br>and 5 minutes<br>before sexual<br>contact in the form<br>of vaginal gel 2.5%<br>oak bark | Not<br>mentioned in<br>the article | Increased satisfaction and orgasm, increased satisfaction from vaginal contraction during intercourse, increased vaginal slipperiness and relaxation after intercourse. | Suberin   |
| Fennel / Nikjo et al.<br>(2018) [16]                               | Fennel   | Clinical<br>trial | 60                         | menopausal<br>women         | Vaginal cream 5<br>grams  | 8 weeks                            | Improving the overall score of sexual performance   | (E)-anethole                                    |
| Date pollen /<br>Yousefzadeh et al.<br>(2017) [17]                 | Date pollen                                      | Clinical<br>trial | 60                         | menopausal<br>women         | 300 mg capsule<br>daily   | 35 days                            | Improvement of orgasm   | dimethoxytoluene<br>isomers                     |
| Elaeagnus<br>angustifolia<br>/Akbarzadeh et al.<br>(2014) [18]     | Elaeagnus<br>angustifolia                        | Clinical<br>trial | 125                        | Non-<br>menopausal<br>women | 4.5 grams in 2 daily<br>doses (capsules<br>every 12 hours)  | 35 days                            | Improvement of orgasm   | 2-<br>Furancarboxaldehyde,<br>5-(hydroxymethyl) |
| Melissa officinalis /<br>Darvish et al. (2018)<br>[20]             | Melissa<br>officinalis                           | Clinical<br>trial | 89                         | Women                       | 500 mg capsule of<br>aqueous extract<br>twice a day   | 4 weeks                            | Improving all areas of sexual performance   | Caffeic acid                                    |
| Ginseng/ Ghorbani<br>et al. (2019) [22]                            | Ginseng  | Clinical<br>trial | 62                         | menopausal<br>women         | 500 mg twice a day  | 4weeks                             | Improving the quality<br>and quantity of sexual<br>performance  | Saponin   |
| Carrot seed/<br>Sadeghi et al. (2020)<br>[23]                      | Carrot seed                                      | Clinical<br>trial | 68                         | women                       | 500 mg capsule 3<br>times a day   | 12 weeks                           | Enhance sexual performance  | carotol   |
| Drimia maritina /<br>Abbasi et al. (2019)<br>[24]                  | Drimia<br>maritina                               | Clinical<br>trial | 60                         | Non-<br>menopausal<br>women | 1-3 times a week<br>and each time 5<br>minutes before<br>intercourse 2-3<br>drops of oil locally                                      | 4 weeks                            | Improving sexual performance in the areas of desire, moisture, orgasm and sexual arousal  | Campesterol                                     |
| Ginkgo biloba/<br>Amiri et al. (2013)<br>[25]                      | Ginkgo biloba                                    | Clinical<br>trial | 77                         | women                       | 60 mg capsule, 120-<br>240 mg daily   | 1 month                            | Enhance sexuality   | 6-hydroxykynurenic<br>acid                      |
| Trigonella foenum<br>graecum /<br>Mazelzadeh et al.<br>(2018) [26] | Trigonella<br>foenum<br>graecum                  | Clinical<br>trial | 60                         | menopausal<br>women         | vaginal cream 5%  | 8 weeks                            | Reducing dyspareunia<br>and increasing libido   | galactomannan                                   |
| Crocus Sativus /<br>Mohammadzadeh et<br>al. (2015) [28]            | Crocus Sativus                                   | Clinical<br>trial | 34                         | Women                       | 30 mg capsules daily  | 4 weeks                            | Enhance sexual performance  | Crocin  |
| Ginger/ Stein et al.<br>(2018) [29]                                | Giner  | Pilot<br>study    | 14                         | Men                         | 100 mg daily  | 30 days                            | Improvement of erectile dysfunction   | Zingiberene                                     |
| Crocus Sativus / Kashani et al. (2013) [27]                        | Crocus Sativus                                   | Clinical<br>trial | 50                         | Men                         | 1% gel before<br>intercourse  | 4 weeks                            | Positive effect on erectile dysfunction   | Crocin  |
| Rosa damascene /<br>Shabanian et al.<br>(2018) [30]                | Rosa<br>damascene and<br>ginger, and<br>cinnamon | Clinical<br>trial | 140                        | Women                       | -   | 2 month                            | Enhance sexual performance and level  | Carboxylic acid                                 |
| Rosa damascene /<br>Fernia et al                                   | Rosa<br>damascene                                | Clinical<br>trial | 50                         | Men                         | drop (oil)  | 4-8                                | Enhance sexual performance and level  | Myrcene   |
| Celery seed /<br>Hasami et al. (2021)<br>[32]                      | Celery seed                                      | Clinical<br>trial | 80                         | men                         | 500 mg capsule 3<br>times a day   | 6 weeks                            | Enhance sexual performance  | Apigenin  |

Table 1: Medicinal plants used in sexual disorders

#### Granatum Punica L.

The results of a study that was conducted on 110 women of reproductive age, after 8 months of intervention with this pomegranate skin gel, it improved the sexual performance of women of reproductive age by enhancing the area of orgasm. The effects of tannin present in pomegranate skin with the effect of appropriate contraction in pelvic floor muscles have caused effective orgasm by increasing the frequency of orgasm and increased sexual satisfaction [14].

#### Ginkgo biloba

Ginkgo biloba extract actually has a significant effect on the two main activities of this plant in the body, including anti-aromatase and vasodilatation, on two important areas of sexual function, i.e. slippery (wetting) and orgasm, and leads to the improvement of sexual function [25, 47].

#### Quercus brantii

It is rich in tannin, which has astringent effects on smooth muscles [48]. In the study of Lorzadeh et al. (2016) which was conducted on Quercus brantii in Lorestan province, the astringent property of it was used and the samples were divided into four groups of 30 people and *Quercus brantii* extract with percentages of 1.5, 2 and 5. 2 was given to them, and the greatest effect was observed in the concentration of 2.5%. In women consuming *Quercus brantii* skin gel, satisfaction and orgasm, feeling of vaginal contraction during intercourse, feeling of wetness (slippery) of the vagina and other parameters including urinary incontinence, feeling of mass coming out of the vagina, feeling of not reaching sexual orgasm, the feeling of air release during intercourse and the feeling of relaxation after intercourse improved in the studied subjects after the intervention [24]. With this mechanism, pelvic floor muscle relaxation causes sagging or prolapse of pelvic organs, dysfunction of pelvic organs, painful intercourse or dyspareunia, back pain, dysfunction and sexual dissatisfaction of couples [49].

#### Lepidium meyenii (Maca)

The effect of *maca* root is to regulate sexual disorders. It is effective on fertility and sexual function in both men and women, as it is mentioned to improve libido and ejaculation, increase the quality, mobility and total number of sperms [50, 51].

#### Foeniculum vulgare (Fennel)

It has phytoestrogenic effects and removes menstrual disorders such as premenstrual syndrome and is effective in treating many diseases [52-54]. *Fennel*, a

plant rich in phytoestrogens, is used in men [58]. In Iranian traditional medicine, fennel is used to treat erectile dysfunction [59]. In traditional Indian medicine, fennel is used for male sexual disorders [60]. This plant exerts its effect on increasing sperm count, testicle weight, and the balance of sex hormones [61].

### Cydonia oblonga

Cydonia oblonga as a strong stimulant, the fruit strengthens the libido. Oral consumption of the hydroalcoholic extract of the plant leads to the improvement of the sexual behavior of the male animal. The results of Najjar et al.'s study (2015) showed that postmenopausal women were divided into two groups of 30 people and the effect of vaginal cream of 5 grams of this plant was compared with placebo. The results indicated the positive effect of this plant compared to placebo, which improved the overall score of sexual function in postmenopausal women based on the FSFI questionnaire [23].

## Panax quinquefolius (Ginseng)

The results of a study conducted on 62 postmenopausal women by receiving 500 mg of this *ginseng* plant twice a day for 28 days, showed the therapeutic effects of this plant on improving the quality of life and the quality and quantity of sexual performance [55]. Researchers have attributed the effect of this plant to the increase in the release of nitric oxide from the vessel wall following the consumption of ginseng, as well as its phytoestrogenic effects, which, by increasing estrogen receptors, has reduced FSH and thus reduced the symptoms of estrogen deficiency [55, 56]. The therapeutic effects of using this plant on the sexual function of men have also been reported [57].

#### Anacyclus pyrethrum

It is a sexual stimulant and can be used to improve sexual function and prevent sexual function disorders and increase sexual desire [62].

#### Ansal (Drimia maritima)

In traditional medicine, *Ansal* has been used for men's sexual disorders, and in the book of Qarabadin Kabir Aghili Khorasani, *Ansal oil* has been very effective in treating erectile dysfunction and increasing libido [63,64].

The results of the study in Iran on 60 women using *Ansal oil* (the oil was prepared from the combination of 900 grams of *Ansal* onion in 3 kilograms of olive oil after heating) for 4 weeks, 1-3 times a week and 5 minutes each time Before intercourse, 2-3 drops of oil topically showed improvement in sexual performance in the areas of desire, moisture, orgasm and sexual excitement [21].

#### Arctium lappa

The results of the animal study show the therapeutic effect of the *Arctium lappa* plant in improving sexual performance and behaviors in a way that increases sexual ability, sexual desire and arousal [65]. Plant extracts and essential oils have important effects on body organs and disease treatment due to the presence of effective medicinal substances, plant bioactive and antioxidants [67-72].

#### Melissa officinalis

The results of a study that was conducted on 89 women aged 18-50 with reduced libido, the treatment group received 500 mg of Melissa officinalis plant extract twice a day for 28 days, the results showed that the use of the plant Compared to placebo, Melissa officinalis (blue extract) improved all areas of women's sexual performance [20].

### Elaeagnus angustifolia

In traditional medicine, it is known as a plant that can have a positive effect on sexual stimulation, especially in women [18]. The results of a clinical trial study that was conducted on 125 women with orgasmic disorders for 35 days showed significant positive effects of this herb compared to placebo on women's orgasmic disorders. Researchers reported that the consumption of this herb improves performance and sexual satisfaction [18].

## Date Palm Pollen

The results of animal studies have shown that date pollen increases testosterone, progesterone, and estradiol in adult male and female mice and as a result improves sexual performance [73, 74].

On the other hand, the presence of compounds such as saponin in date pollen causes the release of nitric oxide and, as a result, improves blood supply to the reproductive system [75]. According to the mentioned mechanisms, using the capsule of this plant in postmenopausal women has improved orgasm compared to the placebo group. The results of a recent study were obtained from the consumption of 300 mg of date pollen for 35 days on 60 postmenopausal women [17].

## *Trifolium pretense* (Red clover)

The results of a study that was conducted on menopausal women, 76 women were studied in the study group, and in the treatment group, a 5-gram applicator was used daily for 8 weeks, and the results showed the therapeutic effects of red clover plant on sexual performance in women. However, in the systematic review study by Najafi et al. (2018), red clover had no effect on the sexual performance of postmenopausal women compared to placebo [15, 76].

#### Lavandula angustifolia

The results of a study showed that the effect of aromatherapy with *Lavandula angustifolia* plant in postmenopausal women reduced anxiety and depression and also improved sexual performance compared to the placebo group [16, 77,78].

#### Pelargonium peltatum (Pelargonium)

The effects of *Pelargonium* aromatherapy on the sexual performance of postmenopausal women have been investigated. In the study of Malkuti et al., which was conducted on 120 postmenopausal women aged 45-55 years, the subjects were randomly assigned to two groups of 60 people consuming Aroma solution and Aroma placebo, which were 2-3 drops 3 times a day. The solution was used by inhalation on the skin of the forearm for 6 weeks. The results of the study showed that combined aromatherapy with *Lavandula*, *Red clover* and *pelargonium* in postmenopausal women led to the improvement of sexual function in all areas except dyspareunia [19].

#### Daucus carota (Carrot seed)

The results of a study showed that 500mg capsules of *carrot seeds* were used 3 times a day for 12 weeks, *carrot seeds* improved the overall sexual performance questionnaire in women [33]. The effects of carrot seeds are attributed to its testosterone-increasing and anti-depressant effects [79-82].

## Apium graveolens (Celery seed)

In traditional medicine, *celery seeds* have been used to treat sexual disorders and affect infertility [83]. In animal studies, the effect of this plant on increasing testosterone levels and its phytoestrogenic effects, which can lead to the improvement of sexual performance, have been reported [84-86]. In the study of Hasami et al., on women, the consumption of 500 mg capsules of this plant led to the improvement of their sexual performance [32]. The results showed that different species of medicinal plants reduce sexual function disorders with different underlying mechanisms.

In fact, a wide range of medicinal plant species with numerous phytochemical compounds such as *Fennel, Celery seed, Trigonella foenum-graecum, Tribulus terrestris, Ros Damascena, Carrot seed, Date pollen, Crocus Sativus, Ginseng, Lavender and Red clover can be mentioned, which are used in the treatment Sexual disorders have been used. Considering the public acceptance and the fact that most people believe that herbal products do not have the side effects of chemical drugs and are in line with their values and lifestyle, it is very important to investigate the mechanisms proposed in the articles in the field of the effects of plants on sexual performance [13]. According to* 

research, medicinal substances in some plants are effective in increasing sexual power by affecting gonads, endocrine system and brain. Herbal medicines with antioxidant effects and anti-inflammatory properties may lead to the improvement of sexual performance of people, such as *Red clover, Crocus Sativus, ginseng and Zingiber officinalis* [12].

There are a number of plants that mimic the function of one or more hormones, hormone mimics have exactly the same physiological effect as the hormone they mimic. These substances do not necessarily need to bind to the receptor for their effect [15]. These substances are used clinically to target the function of testosterone, progesterone and estrogen without binding to their receptors [15].

Some plant species have compounds known as phytoestrogens that mimic the effects of estrogen hormones in the body [14]. Also, some studies have shown that phytoestrogens increase sexual desire and sexual arousal, and by increasing lubrication, it leads to a pleasurable orgasm [16] such as fennel and celery seeds. Herbal treatments that are effective in improving testosterone function not only help to improve sexual performance, but are also able to improve other important functions of testosterone, including increasing adaptation to stressful conditions in both sexes [15] such as Tribulus terrestris, Trigonella foenum-graecum, Red clover and Carrot seeds. Estrogen and testosterone act as key hormones in sexual function, and their decrease causes a decrease in sexual function [13]. An excessive increase in androgens puts these women at risk of insulin resistance, diabetes, dyslipidemia, cancers and cardiovascular diseases.

Herbal anti-androgens (plants inhibiting the activity of excess testosterone) are a class of plant compounds that often reduce tissue sensitivity to androgens or reduce androgenic activity by inhibiting the 5-alpha reductase enzyme, which reduces the conversion of testosterone to androgenic dihydrotestosterones [15]. In addition, at the same time, the reduction of androgens, which play a key role in arousal and sexual desire, affect sexual performance.

Therefore, in people who have a decrease in androgen levels, a plant like *Tribulus terrestris*, increases responses in the area of sexual desire and arousal and improves sexual performance [13]. The mechanism of some plants in improving sexual performance is to reduce anxiety and depression, among these plants we can mention lavender, Red clover and *Zingiber officinalis*. Studies have shown that these plants play a significant role in increasing libido by increasing norepinephrine [17]. The positive effect on the release of nitric oxide is another mechanism of some plants such as *Crocus Sativus* and ginseng in improving sexual

performance. Among the effects of nitric oxide on the cardiovascular system, which is a vasorelaxant factor, it increases blood supply and improves lubrication problems and erectile and ejaculation disorders [13]. Traditional medicine studies provide a unique perspective on the use of medicinal plants in different regions of the world to treat diseases. However, more extensive studies on the mechanism of action of these herbs, their safety and therapeutic effects are needed [84-87].

## Conclusion

Since hormone therapy has many side effects, increasing awareness of how herbs work on androgen, estrogen, and progesterone hormones and the sexual function of people will give us the knowledge that in couples' sexual dysfunctions, there are more choices for correction without using to have hormones with the aim of increasing the quality of life.

## **Author Contributions**

OAA, KDV, YQA, HAK, AH, BMH, HAH hey came up with an idea. HAH, NN, MP literature review. All authors contributed to the article and approved the submission version.

## Conflicts of interest

The authors declare no conflict of interest.

## References

- Goldfarb MR, Trudel G. Marital quality and depression: A review. Marriage & Family Review, )2019(;55(8):737-63.
- Tavakol Z, Nasrabadi AN, Moghadam ZB, Salehiniya H, Rezaei E. A review of the factors associated with marital satisfaction. Galen Medical Journal, (2017);6(3):197-207.
- ghasemi N, rezaei F, Sadeghi M. The Effectiveness of Cognitive-behavioral Therapy on Sexual Function and Quality of Marital Relationships in Women with Sexual Interest/Arousal Disorder. Journal of Nursing Education, (2022);10(3): 87-88.
- Kaliwal, B. B., Parimala, M. D. Dose-Dependent Estrous Cycle, Ovarian Follicles and Biochemical Contents Reversal in Albino Mice after Exposure to Mancozeb. Caspian Journal of Environmental Sciences, (2006); 4(2): 118-131.
- Hayes RD, Bennett CM, Fairley CK, Dennerstein L. What can prevalence studies tell us about female sexual difficulty and dysfunction? The journal of sexual medicine, (2006);3(4):589-95
- Kingsberg SA, Althof S, Simon JA, Bradford A, Bitzer J, Carvalho J, et al. Female sexual dysfunction—medical and psychological treatments, committee 14. The Journal of Sexual Medicine, (2017);14(12):1463-91.
- Basson R, Gilks T. Women's sexual dysfunction associated with psychiatric disorders and their treatment. Women's Health, (2018);14:1745506518762664.
- 8. Schnatz PF, Whitehurst SK, O'Sullivan DM. Sexual dysfunction, depression, and anxiety among patients of an inner-city menopause clinic. Journal of Women's Health, (2010);19(10):1843-9.

## Phytotherapy in sexual disorder: overview of the most important medicinal plants effective on sexual disorders

- Yazdani Karganrood N, Mortazavi P, Motesaddi zarandi S, Eidi A. Effects of inhaled particulate matter (PM2.5) on uterine quality in female rats as an animal model. Caspian Journal of Environmental Sciences, (2023); 1-12.
- Obed AK, Ali AK. Histological and histochemical comparative study on the effects of fasting and a ketogenic diet on the male reproductive system in mice. Caspian Journal of Environmental Sciences, (2023); 1-2.
- Ajao A, Sibiya N, Moteetee A. Sexual prowess from nature: A systematic review of medicinal plants used as aphrodisiacs and sexual dysfunction in sub-Saharan Africa. South African Journal of Botany, (2019);122:342-59.
- Ma A, Mo S. Effect of Medicinal plant of roses on sexual dysfunction of female and male subjects: A systematic review of clinical trials. Journal of Torbat Heydariyeh University of Medical Sciences, (2020);7(4):80-92.
- FARAHMAND M, RAMEZANI TF. The effect of medicinal plants in the treatment of sexual disorders: A narrative Review. Curr Drug Discov Technol, (2021); 18(6): 130921186972.
- Ghavami Z, Gamissi J, Ziaie T, Mollabashi N. Effect of herbal composition (Oak Gall, pomegranate peel and Fenugreek Seeds) on sexual satisfaction of women at reproductive age. The Iranian Journal of Obstetrics, Gynecology and Infertility, (2020); 23(10): 24-33.
- Khayatan J, Iravani M, Moghimipour E, Haghighizadeh MH, Jelodarian P. The effect of red clover vaginal cream on sexual function in postmenopausal women: A randomized, controlled clinical trial. Iran J Obstet Gynecol Infertil, (2019); 22(8):26-34.
- Nikjou R, Kazemzadeh R, Asadzadeh F, Fathi R, Mostafazadeh F. The effect of lavender aromatherapy on the symptoms of menopause. Journal of the National Medical Association, (2018); 110(3):265-9.
- Yosefzadeh S, Sadeghi S, Rakhshandeh H, Dadghar S, Mazloum SR. The effect of date palm pollen capsule on orgasm and sexual satisfaction in menopausal women: a double-blind controlled clinical trial. Iran J Obstet Gynecol Infertil, (2017); 20(10):43-51.
- Akbarzadeh M, Zeinalzadeh S, Zolghadri J, Mohagheghzadeh A, Faridi P, Sayadi M. Comparison of Elaeagnus angustifolia extract and sildenafil citrate on female orgasmic disorders: a randomized clinical trial. Journal of reproduction & infertility, (2014); 15(4):190-8.
- Malakouti J, Farshbaf Khalili A, Asghari Jafarabadi M, Jabbari F. Effect of combined inhaler aromatherapy on sexual function in postmenopausal women: a randomized controlled trial. Iran J Obstet Gynecol Infertil, (2016); 19(1.2):9-15.
- Darvish-Mofrad-Kashani Z, Emaratkar E, Hashem-Dabaghian F, Emadi F, Raisi F, Aliasl J, et al. Effect of Melissa officinalis (lemon balm) on sexual dysfunction in women: A doubleblind, randomized, placebocontrolled study. Iranian Journal of Pharmaceutical Research, (2018); 17(Suppl):89-100.
- Abbasi Pirouz M, Zojaji A, Shakeri MT, Mirzaei K. Effect of Squill on the sexual function among women of reproductive age. Iran J Obstet Gynecol Infertil, (2018); 21(10):57-65.
- Ghorbani Z, Mirghafourvand M, Charandabi SM, Javadzadeh Y. The effect of ginseng on sexual dysfunction in menopausal women: A double-blind, randomized, controlled trial. Complementary Therapies in Medicine, (2019); 1;45:57-64.
- Najar S, Yaralizadeh M, Abedi P, Namjooyan F. Effect of fennel vaginal cream on dysparonia and sexual satisfaction among postmenopausal women: a double-blind randomized controlled trial. Iran J Obstet Gynecol Infertil, (2015); 18(171):8-16.
- Lorzadeh N, Sepavand F, Soleimaninezhad M, Kazemi RN. The effect of extract of internal layer of quercus (OAK GAL) in contraction of vaginal smooth muscles in women with vaginal relaxation. Journal of Medicinal Plants, (2016); 15(60):173-181.

- 25. Amiri PM, Taavoni S, Haghani H. Effect of ginkgo biloba extract on sexual activity in 50–60 years menopausal women in Tehran. Journal of Sabzevar University of Medical Sciences, (2013): 20(2):204-212.
- Mazalzadeh F, Hekmat K, Namjoyan F, SakiMalehi A. Effect of fenugreek vaginal cream on dyspareunia and sexual satisfaction in menopausal women: a randomized clinical trial. Iran J Obstet Gynecol Infertil, (2018); 21(3):22-30.
- 27. Mohammadzadeh-Moghadam H, Nazari SM, Shamsa A, Kamalinejad M, Esmaeeli H, Asadpour AA, et al. Effects of a topical saffron (Crocus sativus L) gel on erectile dysfunction in diabetics: A randomized, parallelgroup, double-blind, placebo-controlled trial. Journal of Evidence-Based Complementary & Alternative Medicine, (2015); 20(4):283-6.
- Kashani L, Raisi F, Saroukhani S, Sohrabi H, Modabbernia A, Nasehi AA, et al. Saffron for treatment of fluoxetine-induced sexual dysfunction in women: randomized double-blind placebo-controlled study. Human Psychopharmacology: Clinical and Experimental, (2013); 28(1):54-60.
- Stein RA, Schmid K, Bolivar J, Swick AG, Joyal SV, Hirsh SP. Kaempferia parviflora ethanol extract improves self-assessed sexual health in men: A pilot study. Journal of Integrative Medicine, (2018); 16(4):249-54.
- Shabanian S, Ebrahimbabaei M, Safavi P, Lotfizadeh M. Comparing the effect of rose drop, ginger, and cinnamon on sexual function in depressed women with sexual dysfunction. Pharmacognosy Research, (2018); 10(3):314.
- 31. Farnia V, Tatari F, Alikhani M, Shakeri J, Taghizadeh M, Karbasizadeh H, et al. Rosa Damascena oil improved sexual function and testosterone in male patients with opium use disorder under methadone maintenance therapy–results from a double-blind, randomized, placebo-controlled clinical trial. Drug and Alcohol Dependence, (2017); 176:117-25.
- Hessami K, Rahnavard T, Hosseinkhani A, Azima S, Sayadi M, Faraji A, et al. Treatment of women's sexual dysfunction using Apium graveolens L. Fruit (celery seed): A double-blind, randomized, placebo-controlled clinical trial. Journal of Ethnopharmacology, (2021); 264:113400.
- 33. Sadeghi S, Bahrami R, Raisi F, Rampisheh Z, Ghobadi A, Akhtari E. Evaluation of the effect of carrot seed (Daucus Carota) in women of fertile age with hypoactive sexual desire disorder: A randomized double-blind clinical trial. Complementary Therapies in Medicine, (2020); 54:102543.
- Ranjbar H, Ashrafizaveh A. Effects of saffron (Crocus sativus) on sexual dysfunction among men and women: A systematic review and meta-analysis. Avicenna Journal of Phytomedicine, (2019); 9(5):419-27.
- Maleki-Saghooni N, Mirzaeii K, Hosseinzadeh H, Sadeghi R, Irani M. A systematic review and meta-analysis of clinical trials on saffron (Crocus sativus) effectiveness and safety on erectile dysfunction and semen parameters. Avicenna Journal of Phytomedicine, (2018); (3):198-209.
- Palacios S, Soler E, Ramírez M, Lilue M, Khorsandi D, Losa F. Effect of a multi-ingredient based food supplement on sexual function in women with low sexual desire. BMC women's Health, (2019); 19(1):1-7.
- 37. Rao A, Steels E, Beccaria G, Inder WJ, Vitetta L. Influence of a specialized Trigonella foenum-graecum seed extract (libifem), on testosterone, estradiol and sexual function in healthy menstruating women, a randomised placebo controlled study. Phytotherapy Research, (2015); 29(8):1123-30.
- 38. Rao A, Steels E, Inder WJ, Abraham S, Vitetta L. Testofen, a specialised Trigonella foenum-graecum seed extract reduces age-related symptoms of androgen decrease, increases testosterone levels and improves sexual function in healthy aging males in a double-blind randomised clinical study. The Aging Male, (2016); 2;19(2):134-42.
- Maheshwari A, Verma N, Swaroop A, Bagchi M, Preuss HG, Tiwari K, Bagchi D. Efficacy of FurosapTM, a novel Trigonella foenum-graecum seed extract, in enhancing testosterone level

## Phytotherapy in sexual disorder: overview of the most important medicinal plants effective on sexual disorders

- and improving sperm profile in male volunteers. International Journal of Medical Sciences, (2017); 14(1): 58.
- Maheshwari A, Verma N, Swaroop A, Bagchi M, Preuss HG, Tiwari K, Bagchi D. Efficacy of FurosapTM, a novel Trigonella foenum-graecum seed extract, in enhancing testosterone level and improving sperm profile in male volunteers. International Journal of Medical Sciences, (2017); 14(1): 58.
- Thompson R, Thompson J, Thompson J, inventors; Js LLC, assignee. Methods to treat one or all of the defined etiologies of female sexual dysfunction. United States patent application US, (2005); 11: 176,546.
- Bombardelli E, Morazzoni P, Riva A, Seghizzi R, inventors; Indena SpA, assignee. Formulations useful in the treatment of male and female impotence. United States patent US, (2008); 7.361.370.
- 43. Lim PH. Asian herbals and aphrodisiacs used for managing ED. Translational Andrology and Urology, (2017); 6(2):167.
- Martimbianco AL, Pacheco RL, Vilarino FL, Latorraca CD, Torloni MR, Riera R. Tribulus Terrestris for Female Sexual Dysfunction: A Systematic Review. Revista Brasileira de Ginecologia e Obstetrícia, (2020); 42(7): 427-35.
- Neychev V, Mitev V. Pro-sexual and androgen enhancing effects of Tribulus terrestris L.: fact or fiction. Journal of Ethnopharmacology, (2016); 17(9): 345-55.
- Akhtari E, Raisi F, Keshavarz M, Hosseini H, Sohrabvand F, Bioos S, Kamalinejad M, Ghobadi A. Tribulus terrestris for treatment of sexual dysfunction in women: randomized double-blind placebo-controlled study. DARU Journal of Pharmaceutical Sciences, (2014); 22(1): 1-7.
- Zhao J, Dasmahapatra AK, Khan SI, Khan IA. Anti-aromatase activity of the constituents from damiana (Turnera diffusa). Journal of Ethnopharmacology, (2008); 120(3): 387-93.
- Haidari R, Siami A, Pakbaz M, Aghazadeh M. Measurement of tannin in four genotype of Quercus infectoria Olive and application of their gall powder in treatment of wound. J Aro Med Pla Res Iran, (2005); 21(4):433-43.
- 49. Artibani W, Haab F, Hilton P. Pelvic floor reconstruction. European Urology, (2002); 42(1):1-6.
- Peres ND, Bortoluzzi LC, Marques LL, Formigoni M, Fuchs RH, Droval AA, et al. Medicinal effects of Peruvian maca (Lepidium meyenii): a review. Food & function, (2020); 11(1):83-92.
- 51. Melnikovova I, Fait T, Kolarova M, Fernandez EC, Milella L. Effect of Lepidium meyenii Walp. on semen parameters and serum hormone levels in healthy adult men: a double-blind, randomized, placebo-controlled pilot study. Evidence-Based Complementary and Alternative Medicine, (2015); 2015: 324369.
- Badgujar SB, Patel VV, Bandivdekar AH. Foeniculum vulgare Mill: a review of its botany, phytochemistry, pharmacology, contemporary application, and toxicology. BioMed research international, 2014; 2014: 842674.
- 53. Akha O, Rabiei K, Kashi Z, Bahar A, Zaeif-Khorasani E, Kosaryan M, et al. The effect of fennel (Foeniculum vulgare) gel 3% in decreasing hair thickness in idiopathic mild to moderate hirsutism, A randomized placebo controlled clinical trial. Caspian Journal of Internal Medicine, (2014); 5(1):26-9.
- 54. Delaram M, Jafari F. The effect of fennel on the pre-menstrual syndrome. Knowledge and Health, (2011); 6(1): 1-6.
- 55. Ghorbani Z, Mirghafourvand M, Charandabi SM, Javadzadeh Y. The effect of ginseng on sexual dysfunction in menopausal women: A double-blind, randomized, controlled trial. Complementary Therapies in Medicine, (2019); 45: 57-64.
- Arring NM, Millstine D, Marks LA, Nail LM. Ginseng as a treatment for fatigue: a systematic review. The Journal of Alternative and Complementary Medicine, (2018); 24(7): 624-
- 57. Park NC, Kim SW, Hwang SY, Park HJ. Efficacy and safety of an herbal formula (KBMSI-2) in the treatment of erectile dysfunction: A preliminary clinical study. Investigative and Clinical Urology, (2019); 60(4): 275-84.

- Haghmorad D, Mahmoudi MB, Haghighi P, Alidadiani P, Shahvazian E, Tavasolian P, et al. Improvement of fertility parameters with Tribulus Terrestris and Anacyclus Pyrethrum treatment in male rats. International Brazilian Journal of Urology, (2019); 45(5):1043-54.
- Ghadiri MK, Gorji A. Natural remedies for impotence in medieval Persia. International Journal of Impotence Research, (2004); 16(1):80-3.
- 60. Ahmed KA, Mitra SK, Venkataraman BV. Assessment of a polyherbal ayurvedic medicine for sexual activity in rats. Indian Drugs, (1999); 36(9): 576-82.
- 61. Sharma V, Boonen J, Spiegeleer BD, Dixit VK. Androgenic and spermatogenic activity of alkylamide-rich e23. Najar S, Yaralizadeh M, Abedi P, Namjooyan F. Effect of fennel vaginal cream on dysparonia and sexual satisfaction among postmenopausal women: a double-blind randomized controlled trial. The Iranian Journal of Obstetrics, Gynecology and Infertility, (2015); 18(171):8-16.
- Aslam M, Sial AA. Effect of hydroalcoholic extract of cydonia oblonga miller (Quince) on sexual behaviour of wistar rats. Advances in Pharmacological Sciences, (2014); 282698.
- 63. Nejatbakhsh F, Karegar-Borzi H, Amin G, Eslaminejad A, Hosseini M, Bozorgi M, et al. Squill Oxymel, a traditional formulation from Drimia Maritima (L.) Stearn, as an add-on treatment in patients with moderate to severe persistent asthma: A pilot, triple-blind, randomized clinical trial. Journal of Ethnopharmacology, (2017): 196:186-92.
- 64. Saadane FZ, Habbachi W, Habbachi S, Boublata NE, Slimani A, Tahraoui A. Toxic effects of Drimia maritima (Asparagaceae) ethanolic extracts on the mortality, development, sexual behaviour and oviposition behaviour of Drosophila melanogaster (Diptera: Drosophilidae). Journal of Animal Behaviour and Biometeorology, (2020); 17;9(1):1
- JianFeng C, PengYing Z, ChengWei X, TaoTao H, YunGui B, KaoShan C. Effect of aqueous extract of Arctium lappa L.(burdock) roots on the sexual behavior of male rats. BMC Complementary and Alternative Medicine, (2012); 12(1):1-8.
- Golkari S, Pourseyedi S, Kazemipour A, Mansouri M. The Effect of Magnetic Iron Oxide Nanoparticles and Ferric Chloride on the Expression of Some Rosmarinic Acid Biosynthetic Genes in Melissa Officinalis L.. Agricultural Biotechnology Journal, (2023); 15(1): 235-254.
- 67. Hosseinnia E, Khara H, Farokhroz M, Yousefi Jourdehi A, Kazemi R. Effects of dietary olive oil and butyl hydroxytoluene (BHT) on digestive and hepatic enzymes, liver and intestinal histology in young Persian sturgeon, Acipenser persicus in brackish water. Aquatic Animals Nutrition, (2020); 6(1): 27-41.
- Ebrahimi Y, Abdalkareem Jasim S, Mohammed BA, A. Salman N, Jabbar AM, M. Hameed N, Goudarzi MA, Parsaei P. Determination of Antioxidant Properties of Mentha longifolia, Pistacia khinjuk and Eucalyptus globulus. Caspian Journal of Environmental Sciences, (2022); 1-6.
- 69. Yeganeh S, Adel M. Effects of dietary rainbow trout visceral protein hydrolysate on hematological, immunological and biochemical serum parameters and antioxidant enzymes of juvenile Siberian sturgeon, Acipenser baerii. Aquatic Animals Nutrition, (2021); 7(4): 43-60.
- Hosseini S M, yousofvand N. Anti-anxiety and antioxidant properties of the zinc nanoparticles green-formulated by an ethnomedicinal plant in the in vivo condition. Plant Biotechnology Persa 2023; 5 (2): URL: http://pbp.medilam.ac.ir/article-1-166-en.html
- Hanifi E, Ahmadifard N, Atashbar B, Meshkini S. Effects of zinc oxide nanoparticles on photosynthetic pigments, zinc accumulation, and activity of antioxidant enzymes of Dunalilla salina. Aquatic Animals Nutrition, (2022); 8(4): 31-
- Shahsavari S, Sarkar S, Sen DJ, Mandal SK. Determination of total antioxidant activity of methanolic extract of Falcaria

#### You're reading

## Phytotherapy in sexual disorder: overview of the most important medicinal plants effective on sexual disorders

- vulgaris. Journal of Biochemicals and Phytomedicine, (2022); 1(1): 8-12.
- Abedi A, Parviz M, Karimian SM, Sadeghipour Rodsari HR. The Effect of Aqueous Extract of Phoenix Dactylifera Pollen Grain on Sexual Behavior of Male Rats. J Phys Pharm Adv, (2012); 2(6):235-42.
- 74. Bahmanpour S, Panjeh SM, Talaei T, Vojdani Z, Poust PA, Zareei S, et al. Effect of Phoenix dactylifera pollen on sperm parameters and reproductive system of adult male rats. Iranian Journal of Medical Sciences, (2006); 31(4):208-212.
- 75. Malini T, Vanithakumari G. Comparative study of the effects of β-sitosterol, estradiol and progesterone on selected biochemical parameters of the uterus of ovariectomised rats. Journal of ethnopharmacology, (1992); 36(1):51-5.
- 76. Najaf Najafi M, Ghazanfarpour M. Effect of phytoestrogens on sexual function in menopausal women: a systematic review and meta-analysis. Climacteric, (2018); 21(5):437-45.
- Almuhsin Ahmed IA, Al-Jboori MJ, AlBahadly ZK. The double infection diagnosis, fungal and insect on date palm offshoot Phoenix dactylifera with treatment pesticides and practical technique. Caspian Journal of Environmental Sciences, (2022); 20(4): 855-858.
- Roozbeh N, Ghazanfarpour M, Khadivzadeh T, Kargarfard L, Dizavandi FR, Shariati K. Effect of Lavender on Sleep, Sexual Desire, Vasomotor, Psychological and Physical Symptom among Menopausal and Elderly Women: A Systematic Review. Journal of Menopausal Medicine, (2019); 25(2):88-93.
- Özliman S, Moradi S, Pirhadi M, Rabeea Banoon S. An Overview of the Most Important Effective Medicinal Plants on Hair Growth and Treatment of Alopecia. Plant Biotechnology Persa, (2021); 3 (2):63-67
- Ahmadi M, Khajoei Nejad F, Dastyar N. Identification of important herbs for anti-fever treatment: an ethnobotanical study of Sojas Rud, northwest Iran. Journal of Biochemicals and Phytomedicine. 2023; 2(2): 54–58.

- 81. Nouri M, Khaki A, Fathiazar F, Rashidi MR. The protective effects of carrot seed extract on spermatogenesis and cauda epididymal sperm reserves in gentamicin treated rats. Cell Journal, (2009); 11(3):327-33.
- 82. Malary M, Khani S, Pourasghar M, Moosazadeh M, Hamzehgardeshi Z. Biopsychosocial determinants of hypoactive sexual desire in women: a narrative review. Materia socio-medica, (2015); 27(6):383-89.
- 83. Al-Asmari AK, Athar MT, Kadasah SG. An updated phytopharmacological review on medicinal plant of Arab region: Apium graveolens linn. Pharmacognosy reviews, (2017); 11(21):13-18.
- 84. Abbasi N, Akhavan MM, Rahbar-Roshandel N, Shafiei M. The effects of low and high concentrations of luteolin on cultured human endothelial cells under normal and glucotoxic conditions: involvement of integrin-linked kinase and cyclooxygenase-2. Phytotherapy Research, (2014);28(9):1301-7.
- 85. Karimi E, Abbasi S, Abbasi N. Thymol polymeric nanoparticle synthesis and its effects on the toxicity of high glucose on OEC cells: involvement of growth factors and integrin-linked kinase. Drug design, Development and Therapy, (2019); 25:2513-32.
- 86. Abbasi N, Ghaneialvar H, Moradi R, Zangeneh MM, Zangeneh A. Formulation and characterization of a novel cutaneous wound healing ointment by silver nanoparticles containing Citrus lemon leaf: A chemobiological study. Arabian Journal of Chemistry, (2021); 1;14(7):103246.
- 87. Bahmani E, Bahmani F, Abbaszadeh S. Ethnobotanical Study of Effective Medicinal Plants as Laxative (Anti-constipation): A Case Study in Arasbaran Forests, Northwest Iran. Journal of Biochemicals and Phytomedicine. 2023; 2(2): 59–63.



This work is licensed under a Creative Commons Attribution-NonCommercial 4.0 International License. To read the copy of this

license please visit:  $\frac{https://creative commons.org/licenses/by-nc/4.0/}{}$