



Full Length Research Article

Advancements in Life Sciences – International Quarterly Journal of Biological Sciences

ARTICLE INFO

Open Access



Date Received:
21/06/2023;
Date Revised:
09/08/2023;
Date Published Online:
20/10/2023;

Staphylococcus aureus isolated from various sites: Detection and diagnosis of some virulence factors production and antibiotic susceptibility

Authors' Affiliation:
Department of Community
Health Technique, Technical
Institute, Middle Technical
University, Baquba - Iraq

Morooj Ali Fahad*, Nabaa Ali Jasim, Shahad Nazar Mustafa, Fatima Malallah Mohammed

***Corresponding Author:**

Morooj Ali Fahad
Email:
Moroojali@mtu.edu.iq

How to Cite:

Fahad MA, Jasim NA,
Mustafa SN, Mohammed FM
(2023). *Staphylococcus aureus* isolated from various
sites: Detection and
diagnosis of some virulence
factors production and
antibiotic susceptibility. Adv.
Life Sci. 10S(1): 46-49.

Keywords:

MRSA; Alpha hemolysin;
Antibiotic

Abstract

Background: Resistance of *Staphylococcus aureus* to commonly used antibiotics is linked to their ability to acquire and disseminate antimicrobial-resistant determinants in nature, and the marine environment may serve as a reservoir for antibiotic-resistant bacteria. Species of this genus can be distinguished by their capability to produce the coagulase enzyme that causes blood clotting, polypeptides that bind to activate prothrombin, in that way converting fibrinogen to fibrin and promoting the clotting of plasma or Blood Fifty skin swabs.

Methods: Samples were collected from patients who are burn and wound infections of Al-Yarmouk teaching hospital, during the period from November 2021 to March 2022, the researchers used different method to diagnosis of antibiotic resistance and alpha hemolysin assay by using a micro titer plate.

Results: The results were found by using cultural traits, microscopic analysis, and biochemical tests, twelve *Staphylococcus aureus* isolates were discovered. A number of antibiotics, including ciprofloxacin/CIP, cefotaxime/CTX, gentamycin/CN, tetracycline/TE, and chloramphenicol/C), were tested using the disc diffusion technique on *Staphylococcus aureus* isolates.

Conclusion: It was discovered that the ten samples had alpha toxin production activity and their susceptibility to antibiotics was known.



Introduction

Staphylococci are facultative anaerobes that are facultatively non-motile and non-spore-forming, and they develop through aerobic respiration. They need a supply of organic nitrogen, such as arginine and valine [1]. The capacity of species in this genus to make polypeptides that bind to activate prothrombin, converting fibrinogen to fibrin, thereby promoting the clotting of plasma or blood, as well as the coagulase enzyme that causes blood clotting, distinguishes them from one another [2]. As a result, *Staphylococcus epidermidis* is a coagulase-negative species, whereas *Staphylococcus aureus* belongs to the coagulase-positive staphylococcal species [3]. A serious threat to global health is the rise of drug-resistant bacterial infections. Many people believe that the widespread use of antibiotics has led to specific forces that have fueled the emergence of resistant strains. Less than a year after the second-generation beta-lactam antibiotics was introduced into clinical practice, Methicillin-resistant *Staphylococcus aureus* (MRSA) was first observed [4]. Due to the way that toxins work and the widespread gene expression of some virulence-determining factors, the present medical treatment methods are in difficulty [5]. Staphylococci species, such as *Staphylococcus aureus* [6]. The bulk of infections can only be treated with methicillin and vancomycin because 90% of *Staphylococcus aureus* strains are penicillin resistant. However, with an increase in reports of methicillin-resistant *Staphylococcus aureus* (MRSA) and vancomycin-resistant enterococci (VRE), chemists are confronted with the challenging task of developing antibiotics with novel modes of action, and physicians with the task of treating infections that appear to have no hope of recovery [7]. because *Staphylococcus aureus* has coagulase production and beta-hemolytic activity, two traits that connect them to virulence in addition to alpha-toxin, which contributes to pathogenesis.

Methods

Isolation and identification of bacteria

From November 2021 to March 2022, fifty skin swabs were taken from patients at Al-Yarmouk Teaching Hospital who had burn and wound infections. Identification of these isolates was also done using the Gram stain and API staph [Biomerieux] as well as the shape, color, size, edges, and height of the colony on the top of brain heart infusion (Himedia) and blood agar dishes [8]. The identification of these isolates also involved the use of a number of assays, such as the catalase test, the mannitol fermentation test, the coagulase test, the alpha-toxin test, and the hemolysis patterns on blood agar and Vitek 2 compact system [9].

Alpha hemolysin assay (Micro titer plate method)

Twelve isolates producing alpha hemolysin toxin were used in this test to check whether it was present in *Staphylococcus aureus* strain supernatants: In Tryptone soya broth, bacteria were grown for 18 hours at 37°C. Each isolate's supernatant was moved to sterile test tubes after centrifugation (7000 rpm) for 15 minutes. 0.1 ml of 2% washed rabbit red blood cells and 0.1 ml of identical bacterial supernatants were added to the well of a U-shaped micro titer plate to test an isolate's capacity to produce alpha hemolysin. Each straight line's first and second wells stand for the negative control, which applied 0.1 ml of ordinary saline and 0.1 ml of washed red blood cells. An hour was spent heating the microtiter plate to 37°C. Plates were maintained at 4°C for 2 to 18 hours following incubation before the outcome (the lysis of RBCs) was determined [10].

Vitek 2 compact system

The completely automated Vitek 2 compact microbiological system uses growth-based technology to identify microorganisms [11]. Colorimetric reagent cards are used in the system, and they are immediately incubated and interpreted. The reagent cards have 64 wells, each of which can hold a different test material. The substrates track a number of metabolic processes, including acidification, alkalization, enzyme hydrolysis, and development in the presence of inhibitors.

Antibiotic sensitivity

The Kirby-Bauer disc diffusion technique was used to test the sensitivity of *Staphylococcus aureus* isolates [12]. During this research, the antibiotic discs ciprofloxacin/CIP, cefotaxime/CTX, gentamycin/CN, tetracycline/TE, and chloramphenicol/C were used.

Results

Morphological characterization

Twelve isolates (24%) were recognized as *Staphylococcus aureus* from the skin swab samples of all isolates. The outcomes showed that these strains were capable of fermenting mannitol and positive for the enzymes catalase, coagulase, and hemolysin (an alpha-toxin). Characterizations based on morphology and biochemistry matched the information provided by [8,9]. In addition to the tests mentioned above, the Api-staph system and the vitek system also perform biochemical identification, which verified the results of the earlier conventional identification shown in the table.

A toxin from *Staphylococcus aureus* called alpha-haemolysin can lyse various cell types and damage membranes. Because alpha-hemolysin disrupts ion transport across host cell membranes, it has several

negative effects on host cells that eventually cause apoptotic cell death and oedema [13]. This is why alpha-hemolysin is believed to be essential in infection. Alpha-hemolysin has been shown to play a part in the virulence of *Staphylococcus aureus* in a variety of infection types, including mastitis and pneumonia. The immunomodulatory characteristics of alpha-haemolysin are also noteworthy, particularly its capacity to promote the release of pro-inflammatory cytokines [14]. As a result, photodynamic therapy that activates alpha-hemolysin may help to both get rid of infectious organisms and defend against damaging inflammatory processes. A number of antibiotics, including ciprofloxacin/CIP, cefotaxime/CTX, gentamycin/CN, tetracycline/TE, and chloramphenicol/C), were tested using the disc diffusion technique on *Staphylococcus aureus* isolates.

Test	Activity
Blood	100%
haemolysis	[+]
Catalase	100%
	[+]
Coagulase	100%
	[+]
Mannitol fermentation	100%
	[+]
Alpha-toxin	100%
Test	[+]

Table 1: Biochemical Tests *S. aureus*.




Figure 1: The percentage of susceptibility of isolates toward the selected antibiotic disks.

Discussion

Humans have been identified as potential contributors of antibiotic-resistant organisms in aquatic settings. They have the ability to transmit antibiotic resistance genes to other diseases and naturally present water bacteria via various mechanisms such as transposons, plasmids, and integrons [15]. Bacterial strains obtained from beach sand, saltwater, and sediments have shown documented resistance to many antimicrobial agents [16].

The prevalence of *Staphylococcus aureus* and MRSA is increasing, leading to a higher frequency of hospital-acquired and community-acquired infections globally. This presents a significant public health issue [17]. In addition, the presence of diverse antibiotics of

industrial origin in aquatic environments has the potential to modify microbial communities [18]. *Staphylococcus aureus* is regarded as a very proficient and adaptable human pathogen owing to its ability to acquire mechanisms of antibiotic resistance and pathogenic determinants, resulting in its development in both healthcare-associated and community contexts [19]. The nosocomial colonization of *Staphylococcus aureus* and methicillin-resistant *Staphylococcus aureus* (MRSA) has the potential to remain unnoticed, with manifestations of the infection becoming apparent many months after patient exposure to the pathogen. Infected individuals might potentially operate as reservoirs for further transmission, particularly due to the prevalence of these strains carrying SCCmec types that encode resistance to methicillin and other beta-lactams [20].

To the best of our knowledge, this is the first study which has used a mixture of phenotypic and genotypic approaches simultaneously to determine the occurrence and antibiotic resistance profiles of *Staphylococcus aureus* strains from beach water and sand in the study area. However, the frequency (12.2%) of isolation was lower in our study than observed in other studies [21]. This study only analyzed a single isolate for every sample, which could account for the lower detection frequency.

The findings of Bimanand's research in 2018 [22], which examined the sensitivity of *Staphylococcus aureus* isolates to Gentamicin and Tetracycline, were in disagreement with those of the present study illustrated in figure 1. Their findings showed that 76% of the isolates were sensitive to Tetracycline and that 69% of the isolates were susceptible to Gentamicin. The findings were consistent with another research on *S. aureus* isolates conducted in India [23], which found that 96% of the isolates were susceptible to chloramphenicol, 76% of the isolates were susceptible to gentamicin, and 8% of the isolates were susceptible to tetracycline. According to the [24] research, about 97.2% of *Staphylococcus aureus* isolates were susceptible to Chloramphenicol, while 61.2% of them were susceptible to Gentamicin. Chloramphenicol showed the highest sensitivity percentage, followed by Gentamicin.

The primary sources and producers of coagulase, catalase, and mannitol fermentation with alpha-hemolysin are skin infections brought on by burns and wounds contaminated with *staphylococcus aureus*. *Staphylococcus aureus*, which was methicillin-resistant was not present in any of the samples. (MRSA). Antibiotics like Ciprofloxacin, Gentamycin, and Chloramphenicol are shown to be sensitive to *Staphylococcus aureus* strains.

Author Contributions

Moroj Ali Fahad: data collection, writing the introduction, objectives, methods, results, and introduction. Nabaa Ali Jasim: data analysis, abstract, keywords, references, and some additions to the discussion, writing the conclusions and make all the corrections and amendments that have been sent by you, Shahad Nazar Mustafa: data collection, and linguistic review. Fatima Malallah Mohammed: reading and reviewing the full article with change or add some sentences.

Competing Interests

The authors declare that they have no conflicts of interest.

References

- Harris G, Foster J, Richards G. An introduction to Staphylococcus aureus and techniques for identifying and quantifying S.aureus adhesins in relation to adhesion to biomaterials: review. *European Cells and Materials*, (2022); 4(1473-2262): 39-60.
- McAdow M, Missiakas D, Schneewind O. Staphylococcus aureus Secretes Coagulase and von Willebrand Factor Binding Protein to Modify the Coagulation Cascade and Establish Host Infections. *Journal of Innate Immunity*, (2012); 4(2): 141-148.
- Ryan K and Ray C. *Sherris Medical Microbiology*, (2004); 4: 370. Publisher McGraw-Hill, New York.
- Harkins C, Pichon B, Doumith M, Parkhill J, Westh H, Tomasz A, de Lencastre H, Bentley S, Kearns A, Holden M. Methicillin-resistant Staphylococcus aureus emerged long before the introduction of methicillin into clinical practice. *Genome biology*, (2017); 18(1): 130.
- Otto M. Staphylococcus aureus toxins. *Current opinion in microbiology*, (2014); 17:32-37.
- Karu T, Tiphlova O, fedoseyeva G. Biostimulating Action of Low-Intensity Monochromatic Visible Light: Is it Possible?. *Laser chemistry*, (1984); 5(0278-6273): 19-25.
- Foster T. *Medical Microbiology*, (1996); Chapter 12. editors Samuel Baron .
- Holt J, Krieg N, Sneath A. *Bergey's manual of determinative bacteriology*, (1996); 9th edition Williams and Wiken.
- Atlas R. *Microorganisms in our world*, (1995); (850): Mosby-your book, Inc.
- Johnson A, Metzger J, Spero L. Production, purification, and chemical characterization of Staphylococcus aureus exfoliative toxin. *infection and immunity*, (1975); 12(5): 1206-1210 .
- Sedaghat H, Nasr E, Mobasherizadeh S, Sallari J, Halaji M, Sadeghi P, Emaneini M, Havaei S. Phenotypic and genotypic characterization of macrolide resistance among Staphylococcus aureus isolates in Isfahan, Iran. *Iran J Microbiol*, (2017); 9(5): 264-270.
- Lombard G, Tealdi S, Lanotte M. The treatment of neurosurgical infections by Laser and prophyryns. *IRIS torino*, (1985); 02A: 364- 366.

- Dinges M, Orwin P, Schlievert P. Exotoxins of Staphylococcus aureus. *Clinical Microbiology*, (2000); 13(1): 16-34.
- Wardenburg J, Bue T, Otto M, Deleo F, Schneewind O. Poring over pores: alpha-hemolysin and panton- Valentine leukociding in Staphylococcus aureus pneumonia. *National Library of Medicine*, (2007); 13(12): (1405-1406).
- Ana O, Paula F, Aline P. Antimicrobial resistance of heterotrophic marine bacteria isolated from seawater and sands of recreational beaches with different organic pollution levels in southeastern Brazil: evidences of resistance dissemination, (2010); 169(1-4): 375-384.
- Manivasagan P, Rajaram G, Ramesh S, Ashokkumar S, Damotharan P. Occurrence and seasonal distribution of antibiotic resistance heterotrophic bacteria and physico-chemical characteristics of Muthupettai mangrove environment, southeast coast of India. *Journal of Environmental Science and Technology*, (2011); 4: 139-149.
- David M and Daum R. Community-Associated Methicillin-Resistant Staphylococcus aureus: Epidemiology and Clinical Consequences of an Emerging Epidemic. *Clinical Microbiology Reviews*, (2010); 23(3): 616-687.
- Zetola N, Francis J, Nuermberger E, Bishai W. Community-acquired methicillin-resistant Staphylococcus aureus: an emerging threat. *Lancet Infectious Diseases*, (2005); 5(5): 275-286.
- Klevens R, Morrison M, Nadle J, Petit S, Gershman K, Ray S, Harrison L, Lynfield R, Dumyati G, Townes J, et al. Invasive methicillin-resistant Staphylococcus aureus infections in the United States. *JAMA*, (2007);298:(1763-1771).
- Boyle-Vavra S and Daum R. Community-acquired methicillin-resistant Staphylococcus aureus: the role of Panton-Valentine leukocidin. *Laboratory Investigation*, (2007); 87(1): 3-9.
- Goodwin K, McNay M, Cao Y, Ebentier D, Madison M, Griffith J. A multi-beach study of Staphylococcus aureus, MRSA, and enterococci in seawater and beach sand. *Water Research*, (2012); 46(13): 4195-207.
- Bimanand L, Taherikalani M, Jalilian F, Sadeghifard N, Ghafourian S, Mahdavi Z, Pakzad I. Association between biofilm production, adhesion genes and drugs resistance in different SCCmec types of methicillin resistant Staphylococcus aureus strains isolated from several major hospitals of Iran. *Iranian journal of basic medical sciences*, (2018); 21(4): 400.
- Ruban W, Babu N, Porteen K, Raja P, Senthilkumar A, Kumarasamy P, Rao A. Prevalence, Antimicrobial susceptibility and Virulence genes of Staphylococcus aureus isolated from pork meat in retail outlets in India. *Journal of Food Safety*, (2018); 39(1): e12589.
- Belbase A, Pant D, Nepal K, Neupane B, Baidhya R, Baidya R, Lekhak B. Antibiotic resistance and biofilm production among the strains of Staphylococcus aureus isolated from pus/wound swab samples in a tertiary care hospital in Nepal. *Annals of Clinical Microbiology and Antimicrobials*, (2017); 16(1):15.



This work is licensed under a Creative Commons Attribution-NonCommercial 4.0 International License. To read the copy of this license please visit: <https://creativecommons.org/licenses/by-nc/4.0/>